

Next-to-Minimal Supersymmetric Standard Model and direct Dark Matter detection

V A Bednyakov^{†1} and H V Klapdor-Kleingrothaus^{‡2}

[†] Laboratory of Nuclear Problems, Joint Institute for Nuclear Research, Moscow region, 141980 Dubna, Russia

[‡] Max-Planck-Institut für Kernphysik, Postfach 103980, D-69029, Heidelberg, Germany

Abstract. Prospects for direct dark matter detection are considered in the Next-to-Minimal Supersymmetric Standard Model (NMSSM). The effective neutralino-quark Lagrangian is obtained and event rates are calculated for ^{73}Ge . Accelerator and cosmological constraints on the NMSSM are included in our numerical analysis, which allows us to conclude that the lightest neutralino from NMSSM could be detected in future dark matter experiments.

1. Introduction

The lightest supersymmetric particle (LSP), the neutralino, is considered now as most promising candidate, which may contribute the main fraction of so-called cold dark matter.

In this paper we investigate direct detection of the relic LSP in the Next-To-Minimal Supersymmetric Standard Model (NMSSM) [1, 2]. The NMSSM is mainly motivated by its potential to eliminate the so-called μ problem [3] of the Minimal Supersymmetric Standard Model (MSSM) [4], where the origin of the μ parameter in the superpotential $W_{\text{MSSM}} = \mu H_1 H_2$ is not understood. For phenomenological reasons it has to be of the order of the electroweak scale, while the "natural" mass scale would be of the order

¹ E-mail: bedny@nusun.jinr.ru

² E-mail: Klapdor@mickey.mpi-hd.mpg.de

of the GUT or Planck scale. This problem is evaded in the NMSSM where the μ term in the superpotential is dynamically generated through $\mu = \lambda x$ with a dimensionless coupling λ and the vacuum expectation value x of the Higgs singlet. Another essential feature of the NMSSM is the fact that the mass bounds for the Higgs bosons and neutralinos are weakened. While in the MSSM experimental data imply a lower mass bound of about 20 GeV for the LSP [5], very light or massless neutralinos and Higgs bosons are not excluded in the NMSSM [6, 7]. The above arguments make an intensive study of the NMSSM phenomenology very desirable. Previously the Higgs and neutralino sectors of the NMSSM were carefully studied in [6]–[11]. The calculation of the LSP relic abundance in the NMSSM was performed for the first time in [12] and recently in [13].

The outline of this paper is as follows. In Sec. 2 we describe the Lagrangian of the NMSSM. Section 3 collects formulas relevant for calculation of the event rate for direct dark matter detection in the framework of the NMSSM. In Sec. 4 we discuss the constraints on the NMSSM parameter space which are used in our analysis. In Sec. 5 we describe our numerical procedure and discuss the results obtained. Sec. 6 contains our conclusion.

2. The Lagrangian of the NMSSM

The NMSSM superpotential is [8] ($\varepsilon_{12} = -\varepsilon_{21} = 1$):

$$W = \lambda \varepsilon_{ij} H_1^i H_2^j N - \frac{1}{3} k N^3 + h_u \varepsilon_{ij} \tilde{Q}^i \tilde{U} H_2^j - h_d \varepsilon_{ij} \tilde{Q}^i \tilde{D} H_1^j - h_e \varepsilon_{ij} \tilde{L}^i \tilde{R} H_1^j, \quad (1)$$

where $H_1 = (H_1^0, H^-)$ and $H_2 = (H^+, H_2^0)$ are the SU(2) Higgs doublets with hypercharge $-1/2$ and $1/2$ and N is the Higgs singlet with hypercharge 0. The notation of the fermion doublets and singlets is conventional, generation indices are omitted. The electroweak gauge-symmetry $SU(2)_I \times U(1)_Y$ is spontaneously broken to the electromagnetic gauge-symmetry $U(1)_{\text{em}}$ by the Higgs VEVs $\langle H_i^0 \rangle = v_i$ with $i = 1, 2$ and $\langle N \rangle = x$, where $v = \sqrt{v_1^2 + v_2^2} = 174$ GeV, $\tan \beta = v_2/v_1$.

The most general supersymmetry breaking potential has the form [8]

$$\begin{aligned} V_{\text{soft}} = & m_1^2 |H_1|^2 + m_2^2 |H_2|^2 + m_3^2 |N|^2 \\ & + m_Q^2 |\tilde{Q}|^2 + m_U^2 |\tilde{U}|^2 + m_D^2 |\tilde{D}|^2 + m_L^2 |\tilde{L}|^2 + m_E^2 |\tilde{R}|^2 \\ & - (\lambda A_\lambda \varepsilon_{ij} H_1^i H_2^j N + \text{h.c.}) - \left(\frac{1}{3} k A_k N^3 + \text{h.c.}\right) \\ & + (h_u A_U \varepsilon_{ij} \tilde{Q}^i \tilde{U} H_2^j - h_d A_D \varepsilon_{ij} \tilde{Q}^i \tilde{D} H_1^j - h_e A_E \varepsilon_{ij} \tilde{L}^i \tilde{R} H_1^j + \text{h.c.}) \\ & + \frac{1}{2} M \lambda^a \lambda^a + \frac{1}{2} M' \lambda' \lambda'. \end{aligned} \quad (2)$$

The minimization conditions for the scalar potential $\partial V/\partial v_{1,2} = 0$ and $\partial V/\partial x = 0$ eliminate three parameters of the Higgs sector which are normally chosen to be m_1^2 , m_2^2 and m_3^2 .

Therefore at the weak scale we are left with the following free parameters of the NMSSM. There are the ratio of the doublet vacuum expectation values, $\tan \beta$, the singlet vacuum expectation value x , the couplings in the superpotential λ and k , the parameters A_λ , A_k , as well as A_U , A_D , A_E (for three generations) in the supersymmetry breaking potential, the gaugino mass parameters M and M' , and the scalar mass parameters for the squarks $m_{Q,U,D}$ and sleptons $m_{L,E}$.

Assuming CP conservation in the Higgs sector, the Higgs CP-even and CP-odd 3×3 matrices are diagonalized by the real orthogonal 3×3 matrices U^S and U^P , respectively,

$$\begin{aligned} \text{Diag}(m_{S_1}^2, m_{S_2}^2, m_{S_3}^2) &= U^{ST} \mathcal{M}_S^2 U^S, \\ \text{Diag}(m_{P_1}^2, m_{P_2}^2, 0) &= U^{PT} \mathcal{M}_P^2 U^P, \end{aligned}$$

where $m_{S_1} < m_{S_2} < m_{S_3}$ and $m_{P_1} < m_{P_2}$ denote the masses of the mass eigenstates of the neutral scalar Higgs bosons S_a ($a = 1, 2, 3$) and neutral pseudoscalar Higgs bosons P_α ($\alpha = 1, 2$) [8]. The Higgs sector of the NMSSM contains five physical neutral Higgs bosons, three Higgs scalars, two pseudoscalars, and two degenerate physical charged Higgs particles $C^\pm$. In our numerical analysis we have included 1-loop radiative corrections to all Higgs mass matrices in accordance with [10, 14].

The masses and mixings of the neutralinos in the basis $\Psi^T = (-i\lambda_1, -i\lambda_2^3, \Psi_{H_1}^0, \Psi_{H_2}^0, \Psi_N)$ are determined by the Lagrangian [8]

$$\mathcal{L} = -\frac{1}{2} \Psi^T M \Psi + \text{h.c.}.$$

The mass of the neutralinos is obtained by diagonalizing the 5×5 matrix M with the orthogonal matrix N :

$$\mathcal{L} = -\frac{1}{2} m_i \bar{\tilde{\chi}}_i^0 \tilde{\chi}_i^0, \quad \tilde{\chi}_i^0 = \begin{pmatrix} \chi_i^0 \\ \bar{\chi}_i^0 \end{pmatrix},$$

with $\chi_i^0 = \mathcal{N}_{ij} \Psi_j$ and $\text{diag}\{m_1, m_2, \dots, m_5\} = N M N^T$. The neutralinos $\tilde{\chi}_i^0$ ($i = 1-5$) are ordered with increasing mass $|m_i|$, thus $\chi \equiv \tilde{\chi}_1^0$ is the LSP neutralino. The matrix elements $\mathcal{N}_{ij}$ ($i, j = 1-5$) describe the composition of the neutralino $\tilde{\chi}_i^0$ in the basis Ψ_j . For example the bino fraction of the lightest neutralino is given by $\mathcal{N}_{11}^2$ and the singlino fraction of this neutralino by $\mathcal{N}_{15}^2$. The remaining particle content is identical with that of the MSSM.

3. Neutralino-nucleus elastic scattering

A dark matter event is elastic scattering of a DM neutralino from a target nucleus producing a nuclear recoil which can be detected by a suitable detector. The corresponding event rate depends on the distribution of the DM neutralinos in the solar vicinity and the cross section of neutralino-nucleus elastic scattering.

An experimentally observable quantity is the differential event rate per unit mass of the target material

$$\frac{dR}{dE_r} = \left[N \frac{\rho_\chi}{m_\chi} \right] \int_{v_{\min}}^{v_{\max}} dv f(v) v \frac{d\sigma}{dq^2}(v, E_r), \quad q^2 = 2M_A E_r.$$

Here $f(v)$ is the velocity distribution of neutralinos in the earth's frame which is usually assumed to be a Maxwellian distribution in the galactic frame. $v_{\max} = v_{\text{esc}} \approx 600$ km/s and $\rho_\chi = 0.3$ GeV·cm⁻³ are the escape velocity and the mass density of the relic neutralinos in the solar vicinity; $v_{\min} = (M_A E_r / 2M_{\text{red}}^2)^{1/2}$ with M_A and M_{red} being the mass of nucleus A and the reduced mass of the neutralino-nucleus system, respectively.

The differential event rate is the most appropriate quantity for comparing with the observed recoil spectrum and allows one to take properly into account spectral characteristics of a specific detector and to separate the background. However, in many cases the total event rate R integrated over the whole kinematical domain of the recoil energy is sufficient. It is widely employed in theoretical papers [15]–[19], [21, 28], for estimating the prospects for DM detection, ignoring experimental complications which may occur on the way. In the present paper we use the total event rate R , because we are going to perform a general analysis aimed at searching for domains with large values of the event rate R .

A general representation of the differential cross section of neutralino-nucleus scattering can be given in terms of three spin-dependent $\mathcal{F}_{ij}(q^2)$ and one spin-independent $\mathcal{F}_S(q^2)$ form factors as follows [20]

$$\begin{aligned} \frac{d\sigma}{dq^2}(v, q^2) = & \frac{8G_F}{v^2} \left(a_0^2 \cdot \mathcal{F}_{00}^2(q^2) + a_0 a_1 \cdot \mathcal{F}_{10}^2(q^2) \right. \\ & \left. + a_1^2 \cdot \mathcal{F}_{11}^2(q^2) + c_0^2 \cdot A^2 \mathcal{F}_S^2(q^2) \right). \end{aligned}$$

The last term corresponding to the spin-independent scalar interaction gains coherent enhancement A^2 (A is the atomic weight of the nucleus in the reaction). The coefficients $a_{0,1}, c_0$ do not depend on nuclear structure but relate to the nucleon structure and the parameters $\mathcal{A}_q, \mathcal{C}_q$ of the relevant low-energy effective neutralino-quark Lagrangian [15]–[18]

$$L_{\text{eff}} = \sum_q \left(\mathcal{A}_q \cdot \bar{\chi} \gamma_\mu \gamma_5 \chi \cdot \bar{q} \gamma^\mu \gamma_5 q + \frac{m_q}{M_W} \cdot \mathcal{C}_q \cdot \bar{\chi} \chi \cdot \bar{q} q \right), \quad (3)$$

where terms with vector and pseudoscalar quark currents are omitted being negligible in the case of non-relativistic DM neutralinos with typical velocities $v \approx 10^{-3}c$. The coefficients in Lagrangian (3) have the form:

$$\begin{aligned}
\mathcal{A}_q = & - \frac{g_2^2}{4M_W^2} \left[\frac{\mathcal{N}_{14}^2 - \mathcal{N}_{13}^2}{2} T_3 \right. \\
& - \frac{M_W^2}{m_{\tilde{q}1}^2 - (m_\chi + m_q)^2} (\cos^2 \theta_q \phi_{qL}^2 + \sin^2 \theta_q \phi_{qR}^2) \\
& - \frac{M_W^2}{m_{\tilde{q}2}^2 - (m_\chi + m_q)^2} (\sin^2 \theta_q \phi_{qL}^2 + \cos^2 \theta_q \phi_{qR}^2) \\
& - \frac{m_q^2}{4} P_q^2 \left(\frac{1}{m_{\tilde{q}1}^2 - (m_\chi + m_q)^2} + \frac{1}{m_{\tilde{q}2}^2 - (m_\chi + m_q)^2} \right) \\
& - \frac{m_q}{2} M_W P_q \sin 2\theta_q T_3 (\mathcal{N}_{12} - \tan \theta_W \mathcal{N}_{11}) \\
& \left. \times \left(\frac{1}{m_{\tilde{q}1}^2 - (m_\chi + m_q)^2} - \frac{1}{m_{\tilde{q}2}^2 - (m_\chi + m_q)^2} \right) \right], \\
\mathcal{C}_q = & - \frac{g_2^2}{4} \left[- \sum_{a=1,2,3} Q_{a11}^{L''} \frac{1}{m_a^2} \left[\left(\frac{1}{2} + T_{3q} \right) \frac{U_{a2}^S}{\sin \beta} + \left(\frac{1}{2} - T_{3q} \right) \frac{U_{a1}^S}{\cos \beta} \right] \right. \\
& + P_q \left(\frac{\cos^2 \theta_q \phi_{qL} - \sin^2 \theta_q \phi_{qR}}{m_{\tilde{q}1}^2 - (m_\chi + m_q)^2} - \frac{\cos^2 \theta_q \phi_{qR} - \sin^2 \theta_q \phi_{qL}}{m_{\tilde{q}2}^2 - (m_\chi + m_q)^2} \right) \\
& + \sin 2\theta_q \left(\frac{m_q}{4M_W} P_q^2 - \frac{M_W}{m_q} \phi_{qL} \phi_{qR} \right) \\
& \left. \times \left(\frac{1}{m_{\tilde{q}1}^2 - (m_\chi + m_q)^2} - \frac{1}{m_{\tilde{q}2}^2 - (m_\chi + m_q)^2} \right) \right].
\end{aligned}$$

Here

$$\begin{aligned}
Q_{a11}^{L''} &= (\mathcal{N}_{12} - \tan \theta_W \mathcal{N}_{11}) [U_{a1}^S \mathcal{N}_{13} - U_{a2}^S \mathcal{N}_{14}] \\
&+ \sqrt{2} \lambda \mathcal{N}_{15} [U_{a1}^S \mathcal{N}_{14} + U_{a2}^S \mathcal{N}_{13}] - 2\sqrt{2} k U_{a3}^S \mathcal{N}_{15}^2, \\
\phi_{qL} &= \mathcal{N}_{12} T_3 + \mathcal{N}_{11} (Q - T_3) \tan \theta_W, \\
\phi_{qR} &= \tan \theta_W Q \mathcal{N}_{11}, \\
P_q &= \left(\frac{1}{2} + T_3 \right) \frac{\mathcal{N}_{14}}{\sin \beta} + \left(\frac{1}{2} - T_3 \right) \frac{\mathcal{N}_{13}}{\cos \beta}.
\end{aligned}$$

The coefficients $\mathcal{A}_q$ and $\mathcal{C}_q$ take into account squark mixing $\tilde{q}_L - \tilde{q}_R$ and the contributions of all CP-even Higgs bosons. Under the assumption $\lambda = k = 0$ these formulas coincide with the relevant formulas in the MSSM [16]. In what follows we use notations and definitions of our papers [19].

4. Constraints on the NMSSM parameter space

Assuming that the neutralinos form a dominant part of the DM in the universe one obtains a cosmological constraint on the neutralino relic density. The present lifetime of the universe is at least 10^{10} years, which implies an upper limit on the expansion rate and correspondingly on the total relic abundance. The contribution of each relic particle species χ has to obey [22]: $\Omega_\chi h_0^2 < 1$, where the relic density parameter $\Omega_\chi = \rho_\chi/\rho_c$ is the ratio of the relic neutralino mass density ρ_χ to the critical one $\rho_c = 1.88 \cdot 10^{-29} h_0^2 \text{ g}\cdot\text{cm}^{-3}$.

To have the parameter $\Omega_\chi h_0^2$ under control we calculate it following the standard procedure on the basis of the approximate formula [23]:

$$\Omega_\chi h_0^2 = 2.13 \times 10^{-11} \left(\frac{T_\chi}{T_\gamma} \right)^3 \left(\frac{T_\gamma}{2.7 K} \right)^3 \times N_F^{1/2} \left(\frac{\text{GeV}^{-2}}{ax_F + bx_F^2/2} \right).$$

Here T_γ is the present day photon temperature, T_χ/T_γ is the reheating factor, $x_F = T_F/m_\chi \approx 1/20$, T_F is the neutralino freeze-out temperature, and N_F is the total number of degrees of freedom at T_F . The coefficients a, b are determined from the non-relativistic expansion $\langle \sigma_{ann} v \rangle \approx a + bx$ of the thermally averaged cross section of neutralino annihilation in the NMSSM. We adopt an approximate treatment not taking into account complications, which occur when the expansion fails [24]. We take into account all possible channels of the χ - χ annihilation in the NMSSM [13].

Since the neutralinos are mixtures of gauginos, higgsinos, and singlino the annihilation can occur both, via s-channel exchange of the Z^0 and Higgs bosons and t-channel exchange of a scalar particle, like a selectron. This constrains the parameter space, as discussed by many groups [23, 25].

In the analysis we ignore possible rescaling of the local neutralino density ρ which may occur in the region of the NMSSM parameter space where $\Omega_\chi h_0^2 < 0.025$ [26]–[28]. If the neutralino is accepted as a dominant part of the DM its density has to exceed the quoted limiting value 0.025. Otherwise the presence of additional DM components should be taken into account and the SUSY solution of the DM problem with such low neutralino density becomes questionable. We assume neutralinos to be a dominant component of the DM halo of our galaxy with a density $\rho_\chi = 0.3 \text{ GeV}\cdot\text{cm}^{-3}$ in the solar vicinity and disregard in the analysis points with $\Omega_\chi h_0^2 < 0.025$.

The parameter space of the NMSSM is constrained by the non-observation of the supersymmetry particles with high energy colliders LEP and Tevatron. Due to the possible presence of singlino component, the $Z\tilde{\chi}^0\tilde{\chi}^0$ and ZS_aP_α couplings can be suppressed in the NMSSM. Therefore the neutralino and Higgs mass bounds are much weaker than in the MSSM [8]. The consequences from the negative neutralino search at LEP for the parameter space and the neutralino masses have been studied in [6, 8].

We used the following constraints from LEP. For new physics contributing to the total Z width $\Delta\Gamma(Z \rightarrow \tilde{\chi}^+ \tilde{\chi}^- + Z \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^0) < 23$ MeV. For new physics contributing to the invisible Z width $\Delta\Gamma(Z \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^0) < 8$ MeV. From the direct neutralino search $B(Z \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_j^0) < 2 \times 10^{-5}$ for $j = 2, \dots, 5$, and $B(Z \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^0) < 5 \times 10^{-5}$, for $i, j = 2, \dots, 5$. We have also included the experimental bounds from the direct search for pseudoscalar Higgs bosons produced together with a Higgs scalar at LEP [5], but this does not significantly affect the excluded parameter domain [8].

In our numerical analysis we use the following experimental restrictions for the SUSY particle spectrum in the NMSSM: $m_{\tilde{\chi}_1^+} \geq 90$ GeV for charginos, $m_{\tilde{\nu}} \geq 80$ GeV for sneutrinos, $m_{\tilde{e}_R} \geq 80$ GeV for selectrons, $m_{\tilde{q}} \geq 150$ GeV for squarks, $m_{\tilde{t}_1} \geq 60$ GeV for light stop, $m_{H^\pm} \geq 65$ GeV for charged Higgses and $m_{S_1} \geq 1$ GeV for the light scalar neutral Higgs. In fact, it appeared that all above-mentioned constraints do not allow m_{S_1} to be smaller than 20 GeV.

5. Numerical Analysis

Randomly scanned parameters of the NMSSM at the Fermi scale are the following: the gaugino mass parameters M' and M , the ratio of the doublet vacuum expectation values, $\tan\beta$, the singlet vacuum expectation value x , the couplings in the superpotential λ and k , squared squark mass parameters $m_{Q_{1,2}}^2$ for the first two generations and $m_{Q_3}^2$ for the third one, the parameters A_λ , A_k , as well as A_t for the third generation. The parameters are varied in the intervals given below

-1000 GeV	$<$	M'	$<$	1000 GeV
-2000 GeV	$<$	M	$<$	2000 GeV
1	$<$	$\tan\beta$	$<$	50
0 GeV	$<$	x	$<$	10000 GeV
-0.87	$<$	λ	$<$	0.87
-0.63	$<$	k	$<$	0.63
100 GeV ²	$<$	$m_{Q_{1,2}}^2$	$<$	1000000 GeV ²
100 GeV ²	$<$	$m_{Q_3}^2$	$<$	1000000 GeV ²
-2000 GeV	$<$	A_t	$<$	2000 GeV
-2000 GeV	$<$	A_λ	$<$	2000 GeV
-2000 GeV	$<$	A_k	$<$	2000 GeV.

For simplicity the other sfermion mass parameters $m_{U_{1,2}}^2$, $m_{D_{1,2}}^2$, $m_{L_{1,2}}^2$, $m_{E_{1,2}}^2$, and m_T^2 , m_B^2 , $m_{L_3}^2$, $m_{E_3}^2$, are chosen to be equal to $m_{Q_2}^2$ and $m_{Q_3}^2$, respectively. Therefore masses of the sfermions in the same generation differ only due to the D-term contribution. Other parameters (except A_t) of the supersymmetry breaking potential A_U , A_D , A_E (for all three generations) are fixed to be zero.




Figure 1. Total event rate R for ^{73}Ge , the LSP gaugino fraction ($\mathcal{N}_{11}^2 + \mathcal{N}_{12}^2$) and singlino fraction ($\mathcal{N}_{15}^2$) versus the LSP mass (from up to down). The left (right) panel presents results obtained without (with) taking into account the cosmological relic density constraint.

The main results of our scan are presented in Figs. 1–4 in the form of scatter plots. Given in Fig. 1 are the total event rate R for ^{73}Ge , and the LSP gaugino fraction ($\mathcal{N}_{11}^2 + \mathcal{N}_{12}^2$) and singlino fraction ($\mathcal{N}_{15}^2$) versus the LSP mass. The left panel in Fig. 1 presents the above-mentioned observables obtained without taking into account the cosmological relic density constraint. In this case the total expected event rate R reaches values up to about 50 events per day and per 1 kg of the ^{73}Ge isotope. As one can see from Fig. 1 the small-mass LSP (less than about 100 GeV) are mostly gauginos, with very small admixture of the singlino component. Large masses of the LSP (larger than 100 GeV) correspond to sizable gaugino and singlino components together with some higgsino fraction. The results of implementation of the cosmological constraint

$$0.025 < \Omega_\chi h_0^2 < 1.$$

one can see in the right panel of Fig. 1. There is approximately a 5-fold reduction of the number of the points which fulfill all restrictions in this case. Nevertheless quite large values of event rate R (above 1 event/day/kg) still survive the cosmological constraint. The gaugino component becomes more significant, but the singlino fraction can not be completely ruled out especially for large masses of the LSP. The higgsino component of the LSP remains still possible only for LSP masses in the vicinity of M_Z . The lower bound for the mass of the LSP now becomes about 5–7 GeV (Fig. 2).




Figure 2. Total event rate R for ^{73}Ge versus the LSP mass in the region of low LSP mass.

Figure 3 shows correlations between mass of the LSP and mass of the lightest scalar Higgs boson S_1 . The upper (about 160 GeV) and lower (about 20 GeV) bound for the mass of S_1 are clear seen. Given in Fig. 4 are the total event rate R for ^{73}Ge versus the masses of all scalar Higgs bosons $S_{1,2,3}$.




Figure 3. Correlations between mass of the LSP and mass of the lightest scalar Higgs boson S_1 .

6. Conclusion

In not too far future new very sensitive dark matter (DM) detectors [29]–[31] may start to operate, and one expects new, very important data from these experiments. The prospects of the direct and indirect detections of the LSP have comprehensively been investigated [15] in the various versions of the MSSM.

In the paper we address the question whether the Next-to-Minimal Supersymmetric Standard Model can be attractive from the point of view of the direct detection of the neutralinos provided the neutralino is the stable LSP. To answer the question we derived the effective low energy neutralino-quark Lagrangian, which takes into account the contributions of extra scalar Higgs boson and extra neutralino. On this basis we calculated the total direct-dark-matter-detection event rate in ^{73}Ge as a representative isotope which is interesting for construction of a realistic dark matter detector. We analyzed the NMSSM taking into account the available accelerator and cosmological constraints by means of random scan of the NMSSM parameter space at the Fermi scale. We demonstrated that the cosmological constraint does not rule out domains in the parameter space which correspond to quite sizable event rate in a germanium detector.




Figure 4. Total event rate R for ^{73}Ge versus the masses of scalar Higgs bosons $S_{1,2,3}$ (from up to down).

Due to relaxation of the gaugino unification condition we found domains in the parameter space where lightest neutralinos have quite small masses (about 7 GeV), acceptable relic abundance and sufficiently large expected event rate for direct detection with a ^{73}Ge -detector.

Therefore the NMSSM looks not worse than the MSSM from the point of view of direct dark matter detection. The question arises: Is it possible to distinguish MSSM and NMSSM by means of direct dark matter detection of LSP? It is a problem to be solved in future. The question can disappear

by itself if negative search for light Higgs with LHC rules out the MSSM. In fact, the upper tree level mass bound for the lightest Higgs scalar of the MSSM

$$m_h^2 \leq m_Z^2 \cos^2 2\beta \quad (4)$$

is increased in the NMSSM to $m_{S_1}^2 \leq m_Z^2 \cos^2 2\beta + \lambda^2(v_1^2 + v_2^2) \sin^2 2\beta$ and the NMSSM may still remain a viable model when the MSSM can be ruled out due to (4). Therefore the NMSSM might remain a viable theoretical background for direct dark matter search for relic neutralinos in the post-MSSM time.

Acknowledgments

We thank S.G. Kovalenko for helpful discussions. The investigation was supported in part (V.A.B.) by Grant GNTP 215 from Russian Ministry of Science and by joint Grant 96-02-00082 from Russian Foundation for Basic Research and Deutsche Forschungsgemeinschaft.

References

- [1] Drees M 1989 *Int. J. of Mod. Phys. A* **4** 3635
- [2] Ellis J, Gunion J F, Haber H E, Roszkowski L and Zwirner F 1989 *Phys. Rev. D* **39** 844
- [3] Kim J E and Nilles H P 1984 *Phys. Lett. B* **138** 150
- [4] Haber H E and Kane G L 1985 *Phys. Rep.* **117** 75
Gunion J F and Haber H E 1986 *Nucl. Phys. B* **272** 1
- [5] Decamp D *et al* 1992 *Phys. Rep.* **216** 253
Adriani O *et al* 1993 *Phys. Rep.* **236** 1
Buskulic D *et al* 1993 *Phys. Lett. B* **313** 312
Akers R *et al* 1994 *Z. Phys. C* **64** 1
- [6] Franke F, Fraas H and Bartl A 1994 *Phys. Lett. B* **336** 415
- [7] Franke F and Fraas H 1995 *Phys. Lett. B* **353** 234
- [8] Franke F and Fraas H 1996 *Z. Phys. C* **72** 309; 1997 *Int. J. Mod. Phys A* **12** (1997) 479
- [9] Ellwanger U 1993 *Phys. Lett. B* **303** 271
- [10] Elliott T, King S F and White P L 1993 *Phys. Lett. B* **314** 56; 1994 *Phys. Rev. D* **49** 2435
- [11] Pandita P N 1993 *Z. Phys. C* **59** 575
- [12] Abel S A, Sarkar S and Whittingham I B 1993 *Nucl. Phys. B* **392** 83
- [13] Stephan A, 1997 *Phys. Lett. B* **411** 97, and hep-ph 9709262.

- [14] King S F and White P L 1995 *Phys. Rev. D* **52** 4183
- [15] Jungman G, Kamionkowski M and Griest K 1996 *Phys. Rep.* **267** 195 and references therein.
- [16] Drees M and Nojiri M M 1993 *Phys. Rev. D* **48** 3483
- [17] Ellis J and Flores R 1991 *Phys. Lett. B* **263** 259; 1993 *Phys. Lett. B* **300** 175; 1993 *Nucl. Phys. B* **400** 2536
- [18] Nath P and Arnowitt R 1997 *Phys. Rev. D* **56** 2820
- [19] Bednyakov V A, Klapdor-Kleingrothaus H V and Kovalenko S G 1994 *Phys. Rev. D* **50** 7128; 1997 *Phys. Rev. D* **55** 503;
Bednyakov V A, Klapdor-Kleingrothaus H V and Kovalenko S G and Ramachers Y 1997 *Z. Phys. A* **357** 339
- [20] Engel J and Vogel P 1989 *Phys. Rev. D* **40** 3132
Engel J, Pitel S and Vogel P 1992 *Int. J. Mod. Phys. E* **1** 1
- [21] Gondolo P and Bergström L 1996 *Astropart. Phys.* **5** 263, hep-ph/9510252.
- [22] Kolb E W and Turner M S 1990 *The early Universe* (Addison-Wesley)
- [23] Drees M and Nojiri M M 1993 *Phys. Rev. D* **47** 376
Ellis J *et al* 1984 *Nucl. Phys. B* **238** 453
- [24] Griest K and Seckel D 1991 *Phys. Rev. D* **43** 3191
Nath P and Arnowitt R 1993 *Phys. Rev. Lett.* **70** 3696
Gelmini G and Gondolo P 1991 *Nucl. Phys. B* **360** 145
- [25] Steigman G *et al* 1986 *Phys. Lett. B* **176** 33
Ellis J *et al* 1986 *Phys. Lett. B* **167** 457
Roszkowski L 1994 *Phys. Rev. D* **50** 4842
Lopez J L, Nanopoulos D V and Pois H 1992 *Phys. Rev. D* **47** 2468
Lopez J L, Nanopoulos D V and Yuan K 1993 *Phys. Rev. D* **48** 2766
Kane G L *et al* 1994 *Phys. Rev. D* **49** 6173
- [26] Gelmini G, Gondolo P and Roulet E 1991 *Nucl. Phys. B* **351** 623
- [27] Bottino A *et al* 1994 *Astropart. Phys.* **2** 77
- [28] Berezhinsky V *et al* 1996 *Astropart. Phys.* **5** 1
- [29] Klapdor-Kleingrothaus H V 1998 *Proc. Int. Conf. "Beyond the Desert"* (Bristol: IOP) p. 485, hep-ex/9802007; 1998 *Yad. Fiz. [Phys. Atomic Nucl.]* **61** 1
Klapdor-Kleingrothaus H V, Hellmig J, and Hirsch M 1998 *J. Phys. G: Nucl. Part. Phys.* **24** 483
- [30] Klapdor-Kleingrothaus H V and Ramachers Y 1998 *Proc. Int. Conf. "Beyond the Desert"* (Bristol IOP) p. 786
- [31] Baudis L *et al* 1997 *Nucl. Instrum. Methods A* **385** 265
Baudis L *et al* 1998 *in this Proceedings*